


AD-A178 196

ESTIMATION OF PARAMETERS IN LATENT CLASS MODELS WITH  
CONSTRAINTS ON THE PARAMETERS (U) PORTLAND STATE UNIV OR 1/1  
DEPT OF PSYCHOLOGY J A PAULSON JUN 86 TR-86-1  
UNCLASSIFIED N00014-81-K-0564 F/G 5/10 NL

END  
DATE  
FBI



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

DTIC



AD-A178 196

ESTIMATION OF PARAMETERS IN  
LATENT CLASS MODELS WITH CONSTRAINTS  
ON THE PARAMETERS

James A. Paulson  
Psychology Department  
Portland State University

June 1986

DTIC  
ELECTE  
MAR 23 1987

This research was sponsored by the Personnel and Training Research Programs, Psychological Sciences Division, Office of Naval Research, under Contract No. N00014-81-K-0564, NR 150-466.

This document has been approved  
for public release and sale; its  
distribution is unlimited.

DTIC FILE COPY

87 3 20 097

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report ONR86-1	2. GOVT ACCESSION NO. AD-A173 174	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Estimation of Parameters in Latent Class Models with Constraints on the Parameters		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) James A. Paulson		8. CONTRACT OR GRANT NUMBER(s) N00014-81-K-0564
9. PERFORMING ORGANIZATION NAME AND ADDRESS Psychology Department Portland State University Portland, OR 97207		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 150-466
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research Arlington, VA 22217		12. REPORT DATE June 1986
		13. NUMBER OF PAGES 32
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Latent class models Marginal maximum likelihood EM algorithm Monotone Homogeneity Unidimensionality Isotonic regression Up-and-down blocks algorithm Item response functions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) → This paper reviews the application of the EM Algorithm to marginal maximum likelihood estimation of parameters in the latent class model and extends the algorithm to the case where there are monotone homogeneity constraints on the item parameters. A likelihood ratio test of the hypothesis of monotone homogeneity is proposed. The hypothesis is of interest because all standard item response theory models assume that it holds. <i>Keywords:</i>		

# ABSTRACT

This paper reviews the application of the EM Algorithm to marginal maximum likelihood estimation of parameters in the latent class model and extends the algorithm to the case where there are monotone homogeneity constraints on the item parameters. A likelihood ratio test of the hypothesis of monotone homogeneity is proposed. The hypothesis is of interest because all standard item response theory models assume that it holds.

NO. 1	✓
NO. 2	
NO. 3	
NO. 4	
NO. 5	
NO. 6	
NO. 7	
NO. 8	
NO. 9	
NO. 10	
NO. 11	
NO. 12	
NO. 13	
NO. 14	
NO. 15	
NO. 16	
NO. 17	
NO. 18	
NO. 19	
NO. 20	
NO. 21	
NO. 22	
NO. 23	
NO. 24	
NO. 25	
NO. 26	
NO. 27	
NO. 28	
NO. 29	
NO. 30	
NO. 31	
NO. 32	
NO. 33	
NO. 34	
NO. 35	
NO. 36	
NO. 37	
NO. 38	
NO. 39	
NO. 40	
NO. 41	
NO. 42	
NO. 43	
NO. 44	
NO. 45	
NO. 46	
NO. 47	
NO. 48	
NO. 49	
NO. 50	
NO. 51	
NO. 52	
NO. 53	
NO. 54	
NO. 55	
NO. 56	
NO. 57	
NO. 58	
NO. 59	
NO. 60	
NO. 61	
NO. 62	
NO. 63	
NO. 64	
NO. 65	
NO. 66	
NO. 67	
NO. 68	
NO. 69	
NO. 70	
NO. 71	
NO. 72	
NO. 73	
NO. 74	
NO. 75	
NO. 76	
NO. 77	
NO. 78	
NO. 79	
NO. 80	
NO. 81	
NO. 82	
NO. 83	
NO. 84	
NO. 85	
NO. 86	
NO. 87	
NO. 88	
NO. 89	
NO. 90	
NO. 91	
NO. 92	
NO. 93	
NO. 94	
NO. 95	
NO. 96	
NO. 97	
NO. 98	
NO. 99	
NO. 100	



A1

## INTRODUCTION

The purpose of this paper is twofold: first, to review the application of the EM algorithm of Dempster, Laird, and Rubin (1977) to marginal maximum likelihood estimation of parameters in the latent class model, and second, to extend this algorithm to the case where there are monotone homogeneity constraints on the item parameters.

Let us briefly review the elements of latent class models. The reader desiring a thorough introduction can consult Lazarsfeld and Henry (1968). The data to be accounted for are vectors of responses to items. In this paper we are only concerned with dichotomous item responses, although many of the ideas can be generalized to the case of polychotomous responses (cf. Goodman, 1974). It is assumed that every subject belongs to exactly one of a finite set of mutually exclusive and exhaustive latent classes. Theoretically, the distribution of the response vectors is to be accounted for by two sets of parameters and one key assumption. The two sets of parameters are the state probabilities,  $(v_k)$ , governing the multinomial distribution of subjects over the latent classes, and the conditional probabilities of correct response to each item, given the respective states,  $(p_{kj})$ . The key assumption is that the responses are conditionally independent, given the state of subject. This implies that any relationships between items must be explained in terms of differences in the  $p_{kj}$ 's between classes. Models are specified by stipulating the number of classes and by placing constraints on the matrix of conditional probabilities.

For some time the problem of estimating parameters in latent class structures presented a real obstacle in the application of the latent class framework. It is necessary to employ iterative procedures in which one selects a set of trial values, improves upon these values in the light of the data via some appropriate algorithm, and then repeats the process until (one hopes) the values stabilize at a good solution. McHugh (1956) derived the maximum likelihood estimators, but his solution applies only to the unconstrained model.

A great advance was achieved when Goodman (1974) described a particularly simple iterative procedure which also has the virtue of automatically producing estimates of probabilities which fall in the unit interval; furthermore it is very easy to modify the procedure to satisfy a fair variety of other constraints on the parameters. There is one problem which Goodman's procedure shares with McHugh's, however. Both procedures take as their data the frequency counts in the cells of the multi-way item-by-item ---by-item contingency table. For a relatively small number of items this presents no problem. But the number of cells grows exponentially with the number of items, so calculations which require dealing with all these cells become impractical very quickly as the number of items is increased. For example, the contingency table for data from a 20 item test would have  $2^{20}$  cells, which is more than a million. Fortunately, it is possible to formulate an algorithm which is logically equivalent to Goodman's, but which circumvents the problem of dealing with all possible cells in the n-way contingency table.

Goodman's algorithm and the modification of it to be presented here are just special cases of the EM algorithm applied to the latent class model. It will be useful to carefully review the rationale behind the application of the EM algorithm in the latent class model context, in order to lay the groundwork for the extension of the algorithm to cover monotone homogeneity constraints on the item parameters. The rationale will be developed in the next section and extended to monotonely homogeneous items in the section after that.

#### APPLICATION OF THE EM ALGORITHM TO THE LATENT CLASS MODEL

Estimation of parameters in the latent class model would be easy if we knew the state of each subject. The maximum likelihood estimates of the distribution of subjects over states would just be the sample proportions falling in the respective states. The estimates of conditional response probabilities to items, given state, would be the corresponding sample proportions of item responses.

The missing data about the states of the respective subjects turns an easy problem into a hard one. Problems with this general character, which would be manageable if only some crucial information were not missing, occur in many contexts. They have inspired numerous special algorithms, often of the following form:



1. Make an initial guess at the parameter values.
2. Using this guess, make an informed guess regarding the missing data.
3. Using this informed guess in place of the missing data, apply the procedure you would ordinarily use to estimate parameter values.
4. Replace the initial guess at the parameter values with the latter estimates and repeat the process until the parameter estimates in steps 1 and 3 no longer differ significantly.

Dempster, Laird, and Rubin (1977) synthesized these many special algorithms into a general approach, which they call the EM algorithm, and showed that under fairly general conditions, if maximum likelihood procedures are used at each iteration in Steps 2 and 3 above, the algorithm converges to marginal maximum likelihood estimates.

In describing how this process works in the case of the latent class model, let us use the following notation.

$$x_{ij} = \begin{cases} 1 & \text{if subject } i \text{ is correct on item } j \\ 0 & \text{if subject } i \text{ is incorrect on item } j; \end{cases}$$

$n$  = the number of subjects;

$J$  = the number of items;

$v_k$  = the probability of a subject being in class or state  $k$ ;

$s$  = the number of latent classes or states;

$\underline{v} = (v_1, \dots, v_s)$ , the vector of state probabilities;

$\underline{x}_i$  = the vector of responses of subject  $i$  to items  $j = 1, \dots, J$ ;

$p_{kj}$  = the conditional probability that a subject in state  $k$  will respond correctly to item  $j$ ;

$P = (p_{kj})$ , the states-by-items matrix of conditional response probabilities;

$\underline{e}_k$  = unit vector with 1 in  $k^{\text{th}}$  coordinate and 0's everywhere else,  
 $k=1, \dots, s$ ;

$\underline{z}_i$  = the unit vector  $\underline{e}_k$  corresponding to the state of subject  $i$ .

Note that  $\underline{e}_k' \underline{z}_i$  is 1 if subject  $i$  is in state  $k$ , and is 0 otherwise.

Let us denote the conditional probability of obtaining response vector  $\underline{x}_i$ , given that subject  $i$  is in state  $k$ , by

$$\begin{aligned} l_k(\underline{x}_i) &= l_k(\underline{x}_i | p, \underline{v}) \\ &= \prod_{j=1}^J p_{kj}^{x_{ij}} (1-p_{kj})^{1-x_{ij}}. \end{aligned} \quad (1)$$

The likelihood of the "complete" data, that is, the joint likelihood of responses and missing state membership vectors  $\underline{z}_i$ , is given by

$$\begin{aligned} L(\underline{x}_1, \dots, \underline{x}_n; \underline{z}_1, \dots, \underline{z}_n | p, \underline{v}) \\ = \prod_{i=1}^n \prod_{k=1}^s [v_k l_k(\underline{x}_i)]^{\underline{e}_k' \underline{z}_i} \end{aligned}$$

Let  $I_k$  be the set of indices for subjects in state  $k$ , i.e. those for whom  $\underline{e}_k' \underline{z}_i = 1$ . Then the likelihood of the complete data can be rewritten as

$$\begin{aligned} L(\underline{x}_1, \dots, \underline{x}_n; \underline{z}_1, \dots, \underline{z}_n | p, \underline{v}) \\ = \prod_{k=1}^s \prod_{i \in I_k} v_k l_k(\underline{x}_i) \end{aligned} \quad (2)$$

$$= \left( \prod_{k=1}^S v_k^{n_k} \right) \left( \prod_{k=1}^S \prod_{j=1}^J p_{kj}^{r_{kj}} (1-p_{kj})^{n_k - r_{kj}} \right) \quad (3)$$

where  $n_k$  denotes the number of subjects in state  $k$  and

$$r_{kj} = \sum_{i \in I_k} x_{ij},$$

denotes the number of correct responses to item  $j$  from subjects in state  $k$ . Thus, the likelihood of the complete data is an exponential family and the set of  $r_{kj}$ 's and  $n_k$ 's are sufficient statistics for the likelihood.

The likelihood function for the complete data is to be contrasted with the marginal likelihood function of the data actually observed. The marginal likelihood of the response vector  $\underline{x}_i$  for a given subject is the average over states of the conditional probabilities of the response vector, given the states,

$$\begin{aligned} l^*(\underline{x}_i) &= l^*(\underline{x}_i | P, \underline{v}) \\ &= \sum_{k=1}^S v_k l_k(\underline{x}_i). \end{aligned} \quad (4)$$

The marginal likelihood of all the observed response vectors is given by

$$\begin{aligned} L^*(\underline{x}_1, \dots, \underline{x}_n | P, \underline{v}) \\ &= E_z \{ L(\underline{x}_1, \dots, \underline{x}_n, \underline{z}_1, \dots, \underline{z}_n | P, \underline{v}) \} \\ &= \prod_{i=1}^n l^*(\underline{x}_i). \end{aligned} \quad (5)$$

It is possible to attack the maximization of Equation 5 directly, but doing so leads to a cumbersome system of nonlinear equations. Another approach suggested by the relationship between the likelihood function for the complete data and the marginal likelihood can be used to maximize the latter indirectly.

Taking logarithms of the likelihood of the complete data in equation 3 yields

$$\log L = \sum_{k=1}^S n_k \log v_k + \sum_{k=1}^S \sum_{j=1}^J [r_{kj} \log p_{kj} + (n_k - r_{kj}) \log(1 - p_{kj})]. \quad (6)$$

If we knew the state of each subject, we could count the  $n_k$ 's and  $r_{kj}$ 's and the standard ratios of these frequencies would be seen to be the maximum likelihood estimators of the  $v_k$ 's and  $p_{kj}$ 's.

Suppose we symbolically calculate the conditional expectation of Equation 6, given the observable response vectors  $\underline{x}_1, \dots, \underline{x}_n$  and trial values of the parameters,  $P_0$  and  $\underline{v}_0$ :

$$\begin{aligned} & E\{\log L(\underline{x}_1, \dots, \underline{x}_n, \underline{z}_1, \dots, \underline{z}_n | P, \underline{v}) | \underline{x}_1, \dots, \underline{x}_n, P_0, \underline{v}_0\} \\ &= \sum_{k=1}^S E_0(n_k) \log \underline{v}_k + \sum_{k=1}^S \sum_{j=1}^J [E_0(r_{kj}) \log p_{kj} + E_0(n_k - r_{kj}) \log(1 - p_{kj})], \end{aligned} \quad (7)$$

where  $E_0(\cdot)$  denotes the conditional expectation, given the responses and trial parameter values,  $E(\cdot | \underline{x}_1, \dots, \underline{x}_n, P_0, \underline{v}_0)$ .

Let  $v_{k0}$  denote the  $k^{\text{th}}$  coordinate of the trial state probability vector  $\underline{v}_0$  and let  $v_{ki}$  denote the conditional probability that subject  $i$

is in state  $k$ , given the subject's responses,  $\underline{x}_i$ , and the trial parameter values  $P_0$  and  $\underline{v}_0$ . By Bayes' theorem,

$$v_{ki} = \frac{v_{k0} l_k(\underline{x}_i | P_0, \underline{v}_0)}{l^*(\underline{x}_i | P_0, \underline{v}_0)} \quad (8)$$

The conditional expectations  $E_0(n_k)$  and  $E_0(r_{kj})$  can be easily computed in terms of the  $v_{ki}$ 's. Recall that the scalar product of the state vector for subject  $i$ ,  $\underline{z}_i$ , and the unit vector corresponding to state  $k$ ,  $\underline{e}_k$ , is 1 if subject  $i$  is in state  $k$  and 0, otherwise. Thus,

$$n_k = \sum_{i=1}^n \underline{e}_k' \underline{z}_i$$

and

$$r_{kj} = \sum_{i=1}^n x_{ij} \underline{e}_k' \underline{z}_i.$$

The expected number of subjects in state  $k$ , given the responses and trial values of the parameters, is

$$\begin{aligned} E_0(n_k) &= E\left(\sum_{i=1}^n \underline{e}_k' \underline{z}_i \mid \underline{x}_1, \dots, \underline{x}_n; P_0, \underline{v}_0\right) \\ &= \sum_{i=1}^n P(\underline{z}_i = \underline{e}_k \mid \underline{x}_i; P_0, \underline{v}_0) \\ &= \sum_{i=1}^n v_{ki}. \end{aligned} \quad (9)$$

The expected number of correct responses to item  $j$  from subjects in state  $k$ , given the responses and trial parameter values, is

$$\begin{aligned} E_0(r_{kj}) &= E\left(\sum_{i=1}^n x_{ij} e_{k-1}^i \mid x_1, \dots, x_n; P_0, \underline{v}_0\right) \\ &= \sum_{i=1}^n x_{ij} P(z_i = e_k \mid x_i, \dots, x; P_0, \underline{v}_0) \\ &= \sum_{i=1}^n x_{ij} v_{ki}. \end{aligned} \tag{10}$$

Equations 8, 9 and 10 enable us to compute numbers to use in Equation 7. Note that the trial values of the parameters  $P_0$  and  $\underline{v}_0$  used to compute the  $E_0(r_{kj})$ 's and  $E_0(p_{kj})$ 's are distinct from parameters  $P$  and  $\underline{v}$  which are free variables in the likelihood functions given in Equations 5, 6 and 7.

It is often relatively easy to maximize Equation 7. If the resulting parameter estimates differ from the trial values  $P_0$  and  $\underline{v}_0$ , they will yield higher values of the marginal likelihood function  $L^*$  than  $P_0$  and  $\underline{v}_0$ , though they may not maximize  $L^*$ . If they do not differ from  $P_0$  and  $\underline{v}_0$ , then  $P_0$  and  $\underline{v}_0$  are also solutions to the marginal likelihood equations which result from setting the partial derivatives of  $\log L^*$  equal to zero. This fact is established by Dempster, et. al (1977) for problems in which the likelihood of the complete data is an exponential family, as is the case here. Sometimes there are multiple possible solutions to the marginal maximum likelihood equations and the question arises whether a given solution is the global maximum of the likelihood function. Ways of dealing with this problem will be discussed later in this section.

Finding values of  $P$  and  $\underline{v}$  to maximize Equation 7 breaks down conveniently into two subproblems: maximization of

$$L_v = \sum_{k=1}^S E_0(n_k) \log v_k \quad (11)$$

with respect to the vector of state probabilities,  $\underline{v}$ , and maximization of

$$L_p = \sum_{k=1}^S \sum_{j=1}^J [E_0(r_{kj}) \log p_{kj} + E_0(n_k - r_{kj}) \log (1 - p_{kj})] \quad (12)$$

with respect to the matrix of conditional response probabilities,  $P$ .

If no constraints are placed on the parameters, the solution to the first problem is given by

$$\begin{aligned} v_k &= \frac{E_0(n_k)}{n} \\ &= \frac{\sum_{i=1}^n v_{ki}}{n} \end{aligned} \quad (13)$$

The solution to the second problem is given by

$$\begin{aligned} p_{kj} &= \frac{E_0(r_{kj})}{E_0(n_k)} \\ &= \frac{\sum_{i=1}^n x_{ij} v_{ki}}{\sum_{i=1}^n v_{ki}} \end{aligned} \quad (14)$$

Let  $\theta$  represent a generic item parameter, possibly affecting several of the  $p_{kj}$ 's. In maximizing the part of Equation 7 which depends on the free item parameters, we set the partial derivative of Equation 11 with respect to  $\theta$  equal to zero; the resulting equation can be arranged to read

$$\sum_{k=1}^J \sum_{j=1}^S v_k' \frac{(p_{kj}' - p_{kj})}{p_{kj}(1-p_{kj})} \cdot \frac{\partial p_{kj}}{\partial \theta} = 0. \quad (15)$$

#### Equality and complementarity constraints

In general, Equation 15 leads to a system of nonlinear equations which can be very difficult to solve. However, there are some special cases which are easy to handle. For example, if there are no constraints on the  $p_{kj}$ 's, then each  $p_{kj}$  is a distinct parameter affecting only one term in the sum in Equation 15. The partial derivative with respect to  $p_{kj}$  itself is 1, all the other partials are 0, and we obtain  $p_{kj}'$  as the solution.

More generally, solution is easy if we only wish to impose equality or complementarity constraints, so that we require  $p_{kj} = \theta$  for one set of  $p_{kj}$ 's,  $p_{kj} = 1-\theta$  for another set, and no  $p_{kj}$  outside of these sets depends on  $\theta$ . Then  $p_{kj}(1-p_{kj})$  equals  $\theta(1-\theta)$ , independent of subscript, for all  $j,k$  such that the partial derivative  $\partial p_{kj} / \partial \theta$  is nonzero. The partial derivative is 1 for  $p_{kj}$ 's equal to  $\theta$  and -1 for those equal to  $1-\theta$ . Let  $I_\theta$  be the set of indices  $j,k$  for which  $p_{kj}$  equals  $\theta$  and  $I_{1-\theta}$  the set for which  $p_{kj}$  is  $1-\theta$ . Then



Equation 14 reduces to a linear equation whose solution is the following weighted combination of  $p'_{kj}$ 's:

$$\theta = \frac{\sum_{j,k \in I_{\theta}} v'_k p'_{kj} + \sum_{j,k \in I_{\bar{\theta}}} v'_k (1-p'_{kj})}{\sum_{j,k \in I_{\theta}} v'_k + \sum_{j,k \in I_{\bar{\theta}}} v'_k} \quad (16)$$

The application of the EM algorithm to estimation of parameters in the latent class model with equality and complementarity constraints can be summarized as follows.

1. Select trial values of the parameters  $P_0$  and  $\underline{v}_0$ .
2. Compute conditional state probabilities for all subjects, using Equation 8.
3. Revise the parameter estimates of  $\underline{v}$  via Equation 13 and the estimates of  $P$  via Equations 14 and 16.
4. Repeat Steps 1 through 3, using the revised estimates as new trial values, until the trial values and the revised values no longer differ significantly.

The key computations in this algorithm involve ratios of counts or estimates of counts in which the denominators are always at least as big as the numerators. The constraint that all estimates lie in the unit interval is therefore automatically satisfied. This is a significant feature of the EM approach not shared by the Newton-Raphson

algorithm when applied to the marginal likelihood function in Equation 5.

The most significant problem which this algorithm is likely to encounter in practice is one that it shares with all existing algorithms that would be practical to use on latent class model problems. It was noted earlier in the paper that the maximum likelihood equations can have multiple solutions. In problems where there are multiple solutions, any iterative algorithm will tend to go to a solution close to the trial values initially selected. The resulting solution may well not be the parameter values that truly maximize the likelihood, particularly if the starting values are selected arbitrarily. It is therefore a good idea to try a variety of plausible sets of starting values.

Goodman (1974) gives an algorithm for estimation of parameters in complex contingency tables where some of the variables are not observable. The specialization of his algorithm to the case of dichotomous responses is essentially equivalent to the algorithm given here. Since it is intended for analysis of contingency tables, it assumes that the joint response data for the subjects is summarized in that form. Latent class model estimation programs implementing Goodman's algorithm, such as Clogg and Sawyer (1981), are limited in terms of the number of items which they can accomodate, because the multi-item contingency table quickly becomes unmanageable as the number of items increases. The form of the algorithm given in this paper

deals with each individual response vector, rather than cell counts in a contingency table. Hence, the effect of increasing the number of items has no effect on the algorithm beyond the increase in running time, which is directly proportional to the number of items. Actually, the effect on running time is more closely proportional to the square of the number of items in applications, such as scaling, in which the number of states in the model is also proportional to the number of items. Nevertheless, the effect is much more manageable than the exponential increase in the number of cells in the contingency table with which an algorithm for analysis of contingency tables must deal.

The EM algorithm in the form presented in this paper can cope with tests comprised of many items, while automatically satisfying the fundamental constraint that the parameter estimates all fall in the unit interval and any further equality and complementarity constraints the investigator may wish to impose on the parameters. This fact, together with computational simplicity at each iteration, makes the algorithm an attractive alternative to other approaches to the calculation of the maximum likelihood estimates for the latent class model. Two questions arise: one about how many models of interest can be formulated using only equality and complementarity constraints, and a second one about the possibility that there are other special kinds of constraints which would also yield easy solutions at each iteration of the algorithm.

The answer to the first question is that many latent class models of interest can be expressed in terms of equality and complementarity

constraints on the parameters, including most of the models which have been proposed to date. The latent distance model of Lazarsfeld and Henry (1968) and the quasi-independence model of Goodman (1975), both of which are generalizations of the Guttman simplex model for scaling response patterns, fall in this category. Dayton and Macready (1976, 1980) have proposed analogs and extensions of these models for applications in the analysis of learning hierarchies; their extensions can also be expressed in terms of equality and complementarity constraints. Paulson (1985) has proposed models for signed-number addition test performance with one latent class for students who have mastered the concept and other latent classes corresponding to classes of subjects exhibiting certain systematic patterns of errors. These models are not scaling models, but they are expressible in terms of equality and complementarity constraints on the parameters.

If only equality and complementarity conditions constrain the item parameters, then each  $p_{kj}$  is influenced by exactly one parameter. This rules out models which characterize each  $p_{kj}$  in terms of conjoint effects of item and state parameters, as the Rasch model does, for example. It also rules out models that impose ordering constraints on the  $p_{kj}$ 's. Thus, while many interesting models can be cast in terms of equality and complementarity constraints, many others cannot. Fortunately, models involving conjoint item and state effects and models imposing ordering constraints can be formulated which lead to easily solved forms of Equation 15, preserving the computational simplicity of the EM algorithm.

### EXTENSION TO MONOTONELY HOMOGENEOUS ITEMS

A set of items is said to be monotonely homogeneous if the probabilities of correct response in different subject states fall in the same order for all items. That is, for every pair of items  $j, j'$  and every pair of states  $k, k'$

$$p_{kj} > p_{k'j} \Rightarrow p_{kj'} \geq p_{k'j'} \quad (16)$$

Any set of items conforming to a unidimensional item response theory which requires the probabilities of correct response to items be monotonically increasing functions of ability is monotonely homogeneous. All standard item response theory models impose this condition. On the other hand, if a set of items is monotonely homogeneous, then the averages of the conditional probabilities of correct response over all the items, given the respective states, must fall in the same order as the conditional probabilities for individual items. Let us define ability level for subjects in a given state to be the average of the conditional probabilities of correct response over all the items, i.e. the "true proportion correct". Consider the function associated with each item which is obtained when one plots the conditional probability of correct response to the item, given state, versus true proportion correct. This function is necessarily monotonically increasing for every item. That is, any monotonely homogeneous set of items is associated with a corresponding set of monotonically increasing item response functions. Thus, monotone

homogeneity of a set of items is a necessary and sufficient condition for the items to be representable by an item response theory model with monotonically increasing item response functions.

The assumption of monotone homogeneity is of interest from a couple of different perspectives. Since it is the minimal assumption concerning the form of the item response function sufficient to yield a model with monotonically increasing tracelines, it is worth considering how far the theory can be developed with no further assumptions regarding the form of the functions. Mokken (1971), who first emphasized the importance of the assumption, Mokken and Lewis (1982), and Lewis (1985) have pursued this idea in developing a nonparametric approach to item response theory. A fundamental problem in this development is the estimation of item response functions. In this section we show how to obtain marginal maximum likelihood estimates for these functions in models restricted to a finite number of states. The restriction to a finite number of ability states would seem to be an extreme limitation on the value of such an approach, but Bock and Aitkin (1981) have shown that it is quite workable in application to standard item response theory models.

From another point of view, the assumption of monotone homogeneity is interesting because it provides a definitive criterion for deciding that a unidimensional representation of responses to a given set of items is inappropriate. Holland (1981) has derived from the assumption a series of necessary conditions observed data must satisfy in order to be capable of representation by a unidimensional item response theory

model. Unlike the assumption itself, these conditions can be tested without estimating the item parameters. The simplest of the conditions is that interitem correlations must be nonnegative. Paulson (1985) has shown that this condition is violated in an analysis of signed-number addition test data from a study by Tatsuka and Birenbaum (1979). Paulson describes a simple latent class model which does give a good account of this data. This model is not a scaling model: the states in the model correspond either to mastery of the concept or to one of a set of systematic misconceptions students fall into regarding the concept. The latter states are not ordered. The nonnegativity of interitem correlations is a simple but weak criterion for testing monotone homogeneity. Holland (1981) gives more stringent tests in terms of nonnegativity of correlations between indices based on combined item responses. We will describe a more direct approach later in this section - the likelihood ratio test of the goodness of fit of the monotonely homogeneous finite-state model compared to the fit of the corresponding latent class model without the monotone homogeneity constraint.

#### Modification of the Algorithm to Provide Monotone Homogeneity

Recall that at each iteration of the EM algorithm, the problem of maximizing the conditional likelihood, given the responses and trial values of the parameters, reduces to maximization of two separate terms, one depending only on the state probability distribution parameters and the other only on item parameters, i.e. parameters

affecting conditional response probabilities, given the subject's state. These terms were given above in Equations 10 and 11.

When the item parameters are unconstrained, each term in the sum in Equation 11 can be maximized separately. If the parameters are constrained, but the constraints apply separately to each item, then the set of terms involving each item can be maximized separately. Monotone homogeneity constraints are of this type: they specify the ordering of the conditional correct response probabilities to a particular item, given the respective states, but say nothing about relationships between response probabilities involving different items.

Thus, the maximization of Equation 11 can be written as

$$\max_p L_p = \sum_{j=1}^J \max_{k=1}^S [E_o(r_{kj}) \log p_{kj} + E_o(n_k - r_{kj}) \log(1 - p_{kj})]. \quad (17)$$

Each of the maximizations on the right hand side of Equation 17 corresponds to the maximum likelihood equation for estimating the success probabilities in  $\underline{s}$  independent groups for a particular item. Carrying out the maximization under ordering constraints has a known solution which bears an interest relation to the algorithm given above for dealing with equality constraints.

Consider the problem of maximum likelihood estimation of proportions in  $\underline{s}$  independent groups. Its solution is the familiar

$$\hat{p}_k = \frac{r_k}{n_k}, \quad \text{for } k=1, \dots, \underline{s}.$$



In our problem,  $r_k$  and  $n_k$  are replaced by  $E_0(r_{kj})$  and  $E_0(n_k)$ , their conditional expectations for item  $j$ , given the observed responses and trial values of the parameters.

Now let us add the constraint that

$$p_1 \leq p_2 \leq \dots \leq p_s .$$

Barlow, et al. (1972) have shown how to treat this problem in terms of isotonic regression. The solution is built upon the unconstrained maximum likelihood estimators just mentioned, which are referred to by Barlow, et al. as basic estimates. These basic estimates are amalgamated for solution blocks of adjacent groups within which each group's estimate is set equal to the weighted average of the  $\hat{p}_k$ 's for the groups comprising the solution block.

The solution blocks are formed as follows. At first each group forms its own block. If the basic estimates for all the groups fall in the right order, then the ordering constraint is not active and the constrained estimates coincide with the basic estimates. A group will continue to form its own solution block unless one or both of the following conditions hold:

- a) its inclusion with the group or adjacent set of groups immediately above it in the hypothesized order would increase the average for the resulting block; or
- b) its inclusion with the group or adjacent set of groups immediately below it in the hypothesized order would decrease the average for the resulting block.

The existence of either condition implies a violation of the ordering constraint which can be remedied by combining the groups involved and setting the estimate of probability correct in each of these groups equal to the weighted average of their basic estimates.

Let the weighted average of the basic estimates in the adjacent set of groups with indices running from  $\underline{t}$  through  $\underline{u}$  be denoted by

$$Av(\underline{t}, \underline{u}) = \frac{\sum_{k=\underline{t}}^{\underline{u}} E_0(n_k) \hat{p}_k}{\sum_{k=\underline{t}}^{\underline{u}} E_0(n_k)} . \quad (18)$$

The constrained maximum likelihood estimates can be expressed in terms of "max-min" formulas in four different but equivalent ways:

$$\begin{aligned} p^*_k &= \max_{\underline{t} < k} \min_{\underline{u} > k} Av(\underline{t}, \underline{u}) \\ &= \min_{\underline{u} > k} \max_{\underline{t} < k} Av(\underline{t}, \underline{u}) \\ &= \max_{\underline{t} < k} \min_{\underline{u} > t} Av(\underline{t}, \underline{u}) \\ &= \min_{\underline{u} > k} \max_{\underline{t} < u} Av(\underline{t}, \underline{u}). \end{aligned} \quad (19)$$

The result given by Equations 18 and 19 is what one would obtain using Equation 16 to impose the constraint that conditional

probabilities of correct response in states belonging to the same solution block must be equal. The main difference between the algorithm to impose simple equality constraints and the algorithm necessary to provide monotone homogeneity is that the solution blocks and the equality constraints implicit in them are not given beforehand and can change from one iteration to the next. The latter algorithm must take this into account.

The Up-and-Down Blocks Algorithm. There are many ways one can determine the solution blocks needed to satisfy Equation 19. Barlow et al. (1972) recommend a procedure due to Kruskal (1964), called the "Up-and-Down Blocks" algorithm. Key terms in the tests used in the algorithm are defined as follows. Let  $B_-$ ,  $B$ ,  $B_+$  be three consecutive blocks in order. Block  $B$  is said to be up-satisfied if  $Av B < Av B_+$ . It is said to be down-satisfied if  $Av B_- < Av B$ . At each stage of the algorithm one block is active; this may be amalgamated with an adjacent block or, if it is up-satisfied and down-satisfied, the next block become active. By convention, the first block in order is down-satisfied and the last block is up-satisfied. The exact sequence of events is as follows.

1. At the start, each state is a separate solution block. State 1 is initially specified to be the active block.
2. Test to see if the active block is up-satisfied. If it is, go to the next step. If it is not, pool the active block with the next higher block; the new block becomes active. Go to Step 3.

3. Test to see if the active block is down-satisfied. If it is, go to Step 4. If it is not, pool the active block with the next lower block; the new block becomes active. Go back to Step 2.
4. If the active block does not contain the highest state, make the next higher block active and go back to Step 2. If the active block contains the highest state, the algorithm is finished.

The sequence of tests and actions to determine the solution blocks is given for a hypothetical example in Table 1. In the example, there are five groups with equal sample sizes, so that unweighted averages are used. For the groups in their hypothesized order, the basic estimates are .50, .60, .70, .40, and .90, respectively. When the algorithm encounters the violation of monotone homogeneity in comparing the third and fourth groups, adjustments are made resulting in the final estimates .50, .57, .57, .57, .90.

Insert Table 1 about here

In summary, monotone homogeneity of items is provided by modifying the EM algorithm for unconstrained marginal maximum likelihood estimation as follows. At each iteration, compute the unconstrained estimates and then apply the Up-and-Down Blocks algorithm to the results for each item. Use these monotonely homogeneous values as trial values on the next iteration. Iterate until the stopping criterion you are using is satisfied.

### A Test for Monotone Homogeneity

When there are  $J$  items on a test and one is fitting an unconstrained latent class model with  $s$  states, there are  $J s$  free item parameters to be estimated. Let  $m_j$  denote the number of level sets determined by the Up-and-Down Blocks algorithm for item  $j$ . The number of free item parameters in the model with the monotone homogeneity constraint is then  $\sum m_j$ . Let  $L_u$  and  $L_m$  denote the maxima of the marginal likelihood function evaluated under the unconstrained and monotonely constrained hypotheses, respectively. If the monotone homogeneity hypothesis is correct, then asymptotically the likelihood ratio test statistic

$$-2 \log \lambda = 2(\log L_u - \log L_m) \quad (20)$$

has a chi-squared distribution with  $J s - \sum m_j$  degrees of freedom. This fact can be used to set up critical regions for tests of the hypothesis.

Example. Figure 1 gives graphs of item response functions for some signed-number addition test data obtained by Tatsuoka and Birenbaum (1979). Five pairs of response functions are depicted - one pair for each of five types of items on the test. Each pair consists of an unconstrained item response function and a function constrained

to be monotonically homogeneous. The analysis refers to a special scoring of responses which only attends to whether the magnitude of the response is correct, disregarding the sign of the answer. The curves given are actually averages of four separate curves, because there were four items of each type. Within types, the curves are practically identical. The types vary in terms of whether the larger of the addends appears first or second in the sum, and in terms of the signs of the addends. An item such as "10+-5" would be of the type designated L+-S on Figure 1, for example.

Tatsuoka and Birenbaum found that if one examines the magnitude of the responses and the sign of the responses to these items separately, some very interesting patterns emerge. Some groups of subjects fall into systematic patterns of errors and correct response which correspond to use of erroneous rules. Paulson (1985) found that a five-state latent class model would give a good account of the magnitude responses. That is why five-state models were used to obtain the curves in Figure 1. Examination of the figures reveals that the unconstrained and monotonically homogeneous curves are very similar for four of the five item types. However, for the type -L+-S, the unconstrained curve is practically "U"-shaped. On the basis of these curves, we would expect to reject the hypothesis of monotone homogeneity. Since there are 20 items on the test and five states in the model, the unconstrained model has 100 free item parameters. It

turns out that the total number of level sets in the monotonically constrained model is 81. Thus, there are 19 degrees of freedom for the chi-squared test. We do in fact reject the null hypothesis:  $\chi^2(19)=82.30$ ,  $p<.0001$ .

Further insight can be obtained by examining the data in Figure 1 from another perspective. Figure 2 shows the profiles of responses to the different types of items for subjects in each of the five states. Subjects in State 4, the next to the highest state in terms of number correct, do well on all item types, except Type -L+-S. Subjects in the lowest state in terms of number correct, State 1, do well on Type -L+-S, but poorly on all the rest. Type -L+-S is the only type on the test for which one should add absolute values of the addends; one should subtract on all the rest. Subjects in State 1 appear to follow the rule, "Always add," whereas subjects in State 4 appear to follow the rule, "Always subtract." Clearly, clusters of subjects following erroneous rules of this sort can lead to violations of monotone homogeneity.

#### SUMMARY

This paper has reviewed the application of the EM algorithm to parameter estimation in the latent class model and shown how it can be used to extend existing algorithms to cover monotone homogeneity constraints on the item parameters. The assumption of monotone homogeneity is interesting from a couple of perspectives. Items on a

test have monotonically increasing item response functions if and only if they are monotonely homogeneous, so the assumption leads to a minimally restrictive form of item response theory. If the assumption is violated, a unidimensional item response theory is clearly inappropriate for the data in question. The paper has shown that, if we restrict ourselves to finite-state latent-class models, we can use the EM algorithm to obtain marginal maximum likelihood estimates of the item response functions under the minimal monotone homogeneity assumption. These "nonparametric" estimates should be very useful when the assumption holds. On the other hand, if the assumption does not hold, we would certainly want to know about it. With the marginal maximum likelihood estimates in hand for both the monotonely homogeneous latent class model and the unconstrained model with the same number of states, we can calculate a direct likelihood ratio test of the monotone homogeneity hypothesis.



REFERENCES

- Barlow, R. E., Bartholomew, D. J., Bremner, J. M., and Brunk, H. D. Statistical inference under order restrictions. New York: John Wiley, 1972.
- Bock, R.D., and Aitkin, M. Marginal maximum likelihood estimation of item parameters: application of an EM algorithm. Psychometrika, 1981, 46, 443-459.
- Clogg, C.C., and Sawyer, D.O. A comparison of alternative models for analyzing the scalability of response patterns. In S. Leinhardt (Ed.), Sociological Methodology 1981. San Francisco: Jossey-Bass, 1981.
- Dayton, C.M., and Macready, G.B. A probabilistic model for validation of behavioral hierarchies. Psychometrika, 1976, 41, 189-204.
- Dayton, C.M., and Macready, G.B. A scaling model with response errors and intrinsically unscalable respondents, Psychometrika, 1980, 45, 343-356.
- Dempster, A.P., Laird, N.M., and Rubin, D.B. Maximum likelihood from incomplete data via the EM algorithm (with Discussion). Journal of the Royal Statistical Society, Series B, 1977, 39, 1-38.
- Goodman, L.A. The analysis of qualitative variables when some of the variables are unobservable. Part I--A modified latent structure approach. American Journal of Sociology, 1974, 79, 1179-1259.
- Goodman, L.A. A new model for scaling response patterns: An application of the quasi-independence concept. Journal of the American Statistical Association, 1975, 70, 755-768.
- Holland, P.W. When are item response models consistent with observed data? Psychometrika, 1981, 46, 79-91.
- Kruskal, J. B. Nonmetric multidimensional scaling: a numerical method. Psychometrika, 1964, 29, 115-129.
- Lazarsfeld, P.F., and Henry, N.W. Latent structure analysis. Boston: Houghton-Mifflin, 1968.

References continued

- Lewis, C. Developments in nonparametric ability estimation. In Weiss, D. J. (ed.) Proceedings of the 1982 Item Response Theory and Computerized Adaptive Testing Conference, Department of Psychology, University of Minnesota, Minneapolis, MN, April 1985.
- McHugh, R.B. Efficient estimation and local identification in latent class analysis. Psychometrika, 1956, 21, 331-347.
- Mokken, R. J. A theory and procedure of scale analysis with applications in political research. New York: deGruyter/Berlin: Mouton: 1971.
- Mokken, R. J. and Lewis, C. A nonparametric approach to the analysis of dichotomous item responses. Applied Psychological Measurement, 1982, 6, 417-430.
- Paulson, J. A. Latent class representation of systematic patterns in test responses. Technical Report ONR85-1, Portland State University, Department of Psychology, May 1985.
- Tatsuoka, K., and Birenbaum, M. The danger of relying solely on diagnostic adaptive testing when prior and subsequent instructional methods are different. Computer-based Education Research Laboratory Report E-S. University of Illinois, 1979.

<u>Group Estimates</u>	<u>Up Satisfied</u>	<u>Down Satisfied</u>	<u>Last Block?</u>	<u>Action</u>
<u>.50</u> .60 .70 .40 .90	Yes	Yes	No	Make next block active
.50 <u>.60</u> .70 .40 .90	Yes	Yes	No	Make next block active
.50 .60 <u>.70</u> .40 .90	No			Pool up
.50 .60 <u>.55</u> <u>.55</u> .90		No		Pool down
.50 <u>.57</u> <u>.57</u> <u>.57</u> .90	Yes	Yes	No	Make next block active
.50 .57 .57 .57 <u>.90</u>	Yes	Yes	Yes	Stop.

Table 1. Illustration of the "Up-and-Down Blocks" Algorithm.

Note: Each line indicates the outcomes of test made on an active block. The current estimate of the correct response probabilities for groups comprising the active block are underlined at the left. The action taken is given at the right.




FIGURE 1. Comparison of monotonically homogeneous and unconstrained item response functions.




FIGURE 2. Comparison of monotonically homogeneous and unconstrained state response profiles.

## Portland State University/Paulson

Personnel Analysis Division,  
AF/MPXA  
5C360, The Pentagon  
Washington, DC 20330

Air Force Human Resources Lab  
AFHRL/MPD  
Brooks AFB, TX 78235

Dr. Earl A. Alluisi  
HQ, AFHRL (AFSC)  
Brooks AFB, TX 78235

Dr. Erling B. Andersen  
Department of Statistics  
Studiestraede 6  
1455 Copenhagen  
DENMARK

Dr. Phipps Arabie  
University of Illinois  
Department of Psychology  
603 E. Daniel St.  
Champaign, IL 61820

Technical Director, ARI  
5001 Eisenhower Avenue  
Alexandria, VA 22333

Dr. Eva L. Baker  
UCLA Center for the Study  
of Evaluation  
145 Moore Hall  
University of California  
Los Angeles, CA 90024

Dr. Isaac Bejar  
Educational Testing Service  
Princeton, NJ 08450

Dr. Menucha Birenbaum  
School of Education  
Tel Aviv University  
Tel Aviv, Ramat Aviv 69978  
ISRAEL

Dr. Arthur S. Blaiwes  
Code N711  
Naval Training Equipment Center  
Orlando, FL 32813

Dr. R. Darrell Bock  
University of Chicago  
Department of Education  
Chicago, IL 60637

Cdt. Arnold Bohrer  
Sectie Psychologisch Onderzoek  
Rekruterings-En Selectiecentrum  
Kwartier Koningen Astrid  
Bruijnstraat  
1120 Brussels, BELGIUM

Dr. Robert Breaux  
Code N-095R  
NAVTRAEQUIPCEN  
Orlando, FL 32813

Dr. Robert Brennan  
American College Testing  
Programs  
P. O. Box 168  
Iowa City, IA 52243

Dr. Patricia A. Butler  
NIE Mail Stop 1806  
1200 19th St., NW  
Washington, DC 20208

Mr. James W. Carey  
Commandant (G-PTE)  
U.S. Coast Guard  
2100 Second Street, S.W.  
Washington, DC 20593

Dr. James Carlson  
American College Testing  
Program  
P.O. Box 168  
Iowa City, IA 52243

Dr. John B. Carroll  
409 Elliott Rd.  
Chapel Hill, NC 27514

Dr. Robert Carroll  
NAVOP 01B7  
Washington, DC 20370

Dr. Norman Cliff  
Department of Psychology  
Univ. of So. California  
University Park  
Los Angeles, CA 90007

## Portland State University/Paulson

Director,  
Manpower Support and  
Readiness Program  
Center for Naval Analysis  
2000 North Beauregard Street  
Alexandria, VA 22311

Dr. Stanley Collyer  
Office of Naval Technology  
Code 222  
800 N. Quincy Street  
Arlington, VA 22217-5000

Dr. Hans Crombag  
University of Leyden  
Education Research Center  
Boerhaavelaan 2  
2334 EN Leyden  
The NETHERLANDS

CTR/McGraw-Hill Library  
2500 Garden Road  
Monterey, CA 93940

Dr. Dattprasad Divgi  
Center for Naval Analysis  
4401 Ford Avenue  
P.O. Box 16268  
Alexandria, VA 22302-0268

Dr. Hei-Ki Dong .  
Ball Foundation  
800 Roosevelt Road  
Building C, Suite 206  
Glen Ellyn, IL 60137

Defense Technical  
Information Center  
Cameron Station, Bldg 5  
Alexandria, VA 22314  
Attn: TC  
(12 Copies)

Dr. Stephen Dunbar  
Lindquist Center  
for Measurement  
University of Iowa  
Iowa City, IA 52242

Dr. James A. Earles  
Air Force Human Resources Lab  
Brooks AFB, TX 78235

Dr. Kent Eaton  
Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22333

Dr. John M. Eddins  
University of Illinois  
252 Engineering Research  
Laboratory  
103 South Mathews Street  
Urbana, IL 61801

Dr. Susan Embretson  
University of Kansas  
Psychology Department  
Lawrence, KS 66045

ERIC Facility-Acquisitions  
4833 Rugby Avenue  
Bethesda, MD 20014

Dr. Benjamin A. Fairbank  
Performance Metrics, Inc.  
5825 Callaghan  
Suite 225  
San Antonio, TX 78228

Dr. Leonard Feldt  
Lindquist Center  
for Measurement  
University of Iowa  
Iowa City, IA 52242

Dr. Richard L. Ferguson  
American College Testing  
Program  
P.O. Box 168  
Iowa City, IA 52240

Dr. Gerhard Fischer  
Liebiggasse 5/3  
A 1010 Vienna  
AUSTRIA

Prof. Donald Fitzgerald  
University of New England  
Department of Psychology  
Armidale, New South Wales 2351  
AUSTRALIA

Mr. Paul Foley  
Navy Personnel R&D Center  
San Diego, CA 92152

## Portland State University/Paulson

Dr. Carl H. Frederiksen  
McGill University  
3700 McTavish Street  
Montreal, Quebec H3A 1Y2  
CANADA

Dr. Robert D. Gibbons  
University of Illinois-Chicago  
P.O. Box 6998  
Chicago, IL 60680

Dr. Janice Gifford  
University of Massachusetts  
School of Education  
Amherst, MA 01003

Dr. Robert Glaser  
Learning Research  
& Development Center  
University of Pittsburgh  
2939 O'Hara Street  
Pittsburgh, PA 15260

Dr. Bert Green  
Johns Hopkins University  
Department of Psychology  
Charles & 34th Street  
Baltimore, MD 21218

Dr. Ronald K. Hambleton  
Prof. of Education & Psychology  
University of Massachusetts  
at Amherst  
Hills House  
Amherst, MA 01003

Ms. Rebecca Hetter  
Navy Personnel R&D Center  
Code 62  
San Diego, CA 92152

Dr. Paul W. Holland  
Educational Testing Service  
Rosedale Road  
Princeton, NJ 08541

Prof. Lutz F. Hornke  
Universitat Dusseldorf  
Erziehungswissenschaftliches  
Universitätsstr. 1  
Dusseldorf 1  
WEST GERMANY

Dr. Paul Horst  
677 G Street, #184  
Chula Vista, CA 90010

Mr. Dick Hoshaw  
NAVOP-135  
Arlington Annex  
Room 2834  
Washington, DC 20350

Dr. Lloyd Humphreys  
University of Illinois  
Department of Psychology  
603 East Daniel Street  
Champaign, IL 61820

Dr. Steven Hunka  
Department of Education  
University of Alberta  
Edmonton, Alberta  
CANADA

Dr. Huynh Huynh  
College of Education  
Univ. of South Carolina  
Columbia, SC 29208

Dr. Robert Jannarone  
Department of Psychology  
University of South Carolina  
Columbia, SC 29208

Dr. Douglas H. Jones  
Advanced Statistical  
Technologies Corporation  
16 Trafalgar Court  
Lawrenceville, NJ 08148

Dr. G. Gary Kingsbury  
Portland Public Schools  
Research & Evaluation Department  
501 North Dixon Street  
P. O. Box 3107  
Portland, OR 97209-3107

Dr. William Koch  
University of Texas-Austin  
Measurement and Evaluation  
Center  
Austin, TX 78708



## Portland State University/Paulson

Dr. Leonard Kroeker  
Navy Personnel R&D Center  
San Diego, CA 92152

Dr. Michael Levine  
Educational Psychology  
210 Education Bldg.  
University of Illinois  
Champaign, IL 61801

Dr. Charles Lewis  
Faculteit Sociale Wetenschappen  
Rijksuniversiteit Groningen  
Oude Boteringestraat 23  
9712GC Groningen  
The NETHERLANDS

Dr. Robert Linn  
College of Education  
University of Illinois  
Urbana, IL 61801

Dr. Robert Lockman  
Center for Naval Analysis  
4401 Ford Avenue  
P.O. Box 16268  
Alexandria, VA 22302-0268

Dr. Frederic M. Lord  
Educational Testing Service  
Princeton, NJ 08541

Dr. James Lumsden  
Department of Psychology  
University of Western Australia  
Nedlands W.A. 6009  
AUSTRALIA

Dr. William L. Maloy  
Chief of Naval Education  
and Training  
Naval Air Station  
Pensacola, FL 32508

Dr. Gary Marco  
Stop 21-E  
Educational Testing Service  
Princeton, NJ 08541

Dr. Clessen Martin  
Army Research Institute  
5001 Eisenhower Blvd.  
Alexandria, VA 22333

Dr. James McBride  
Psychological Corporation  
c/o Harcourt, Brace,  
Javanovich Inc.  
1250 West 6th Street  
San Diego, CA 92101

Dr. Clarence McCormick  
HQ, MEPCOM  
MEPCT-P  
2500 Green Bay Road  
North Chicago, IL 60064

Mr. Robert McKinley  
University of Toledo  
Department of Educational Psychology  
Toledo, OH 43606

Dr. Barbara Means  
Human Resources  
Research Organization  
1100 South Washington  
Alexandria, VA 22314

Dr. Robert Mislevy  
Educational Testing Service  
Princeton, NJ 08541

Headquarters, Marine Corps  
Code MPI-20  
Washington, DC 20380

Dr. W. Alan Nicewander  
University of Oklahoma  
Department of Psychology  
Oklahoma City, OK 73069

Dr. William E. Nordbrock  
FMC-ADCO Box 25  
APO, NY 09710

Dr. Melvin R. Novick  
356 Lindquist Center  
for Measurement  
University of Iowa  
Iowa City, IA 52242

Director, Manpower and Personnel  
Laboratory,  
NPRDC (Code 06)  
San Diego, CA 92152

## Portland State University/Paulson

Library, NPRDC  
Code P201L  
San Diego, CA 92152

Commanding Officer,  
Naval Research Laboratory  
Code 2627  
Washington, DC 20390

Dr. James Olson  
WICAT, Inc.  
1875 South State Street  
Orem, UT 84057

Office of Naval Research,  
Code 1142PT  
800 N. Quincy Street  
Arlington, VA 22217-5000  
(6 Copies)

Special Assistant for Marine  
Corps Matters,  
ONR Code 00MC  
800 N. Quincy St.  
Arlington, VA 22217-5000

Dr. Judith Orasanu  
Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22332

Wayne M. Patience  
American Council on Education  
GED Testing Service, Suite 20  
One Dupont Circle, NW  
Washington, DC 20036

Dr. James Paulson  
Department of Psychology  
Portland State University  
P.O. Box 751  
Portland, OR 97207

Dr. Roger Pennell  
Air Force Human Resources  
Laboratory  
Lowry AFB, CO 80230

Dr. Mark D. Reckase  
ACT  
P. O. Box 168  
Iowa City, IA 52242

Dr. Malcolm Ree  
AFHRL/MP  
Brooks AFB, TX 78235

Dr. Carl Ross  
CNET-PDCD  
Building 90  
Great Lakes NTC, IL 60088

Dr. J. Ryan  
Department of Education  
University of South Carolina  
Columbia, SC 29208

Dr. Fumiko Samejima  
Department of Psychology  
University of Tennessee  
Knoxville, TN 37916

Mr. Drew Sands  
NPRDC Code 62  
San Diego, CA 92152

Dr. Robert Sasnor  
Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22332

Dr. Mary Schratz  
Navy Personnel R&D Center  
San Diego, CA 92152

Dr. W. Steve Sellman  
OASD(MRA&L)  
2B269 The Pentagon  
Washington, DC 20301

Dr. Kazuo Shigemasu  
7-9-24 Kugenuma-Kaigan  
Fujisawa 251  
JAPAN

Dr. William Sims  
Center for Naval Analysis  
4401 Ford Avenue  
P.O. Box 16268  
Alexandria, VA 22302-0268

## Portland State University/Paulson

Dr. H. Wallace Sinaiko Manpower Research and Advisory Services Smithsonian Institution 801 North Pitt Street Alexandria, VA 22314	Dr. Maurice Tatsuoka 220 Education Bldg 1310 S. Sixth St. Champaign, IL 61820
Dr. Richard Sorensen Navy Personnel R&D Center San Diego, CA 92152	Dr. David Thissen Department of Psychology University of Kansas Lawrence, KS 66044
Dr. Paul Speckman University of Missouri Department of Statistics Columbia, MO 65201	Mr. Gary Thomasson University of Illinois Educational Psychology Champaign, IL 61820
Dr. Martha Stocking Educational Testing Service Princeton, NJ 08541	Dr. Robert Tsutakawa The Fred Hutchinson Cancer Research Center Division of Public Health Sci. 1124 Columbia Street Seattle, WA 98104
Dr. Peter Stoloff Center for Naval Analysis 200 North Beauregard Street Alexandria, VA 22311	Dr. Ledyard Tucker University of Illinois Department of Psychology 603 E. Daniel Street Champaign, IL 61820
Dr. William Stout University of Illinois Department of Mathematics Urbana, IL 61801	Dr. Vern W. Urry Personnel R&D Center Office of Personnel Management 1900 E. Street, NW Washington, DC 20415
Maj. Bill Strickland AF/MPXOA 4E168 Pentagon Washington, DC 20330	Dr. David Vale Assessment Systems Corp. 2233 University Avenue Suite 310 St. Paul, MN 55114
Dr. Hariharan Swaminathan Laboratory of Psychometric and Evaluation Research School of Education University of Massachusetts Amherst, MA 01003	Dr. Frank Vicino Navy Personnel R&D Center San Diego, CA 92152
Mr. Brad Sympson Navy Personnel R&D Center San Diego, CA 92152	Dr. Howard Wainer Division of Psychological Studies Educational Testing Service Princeton, NJ 08541
Dr. Kikumi Tatsuoka CERL 252 Engineering Research Laboratory Urbana, IL 61801	

## Portland State University/Paulson

Dr. Ming-Mei Wang  
Lindquist Center  
for Measurement  
University of Iowa  
Iowa City, IA 52242

Mr. Thomas A. Warm  
Coast Guard Institute  
P. O. Substation 18  
Oklahoma City, OK 73169

Dr. Brian Waters  
Program Manager  
Manpower Analysis Program  
HumRRO  
1100 S. Washington St.  
Alexandria, VA 22314

Dr. David J. Weiss  
N660 Elliott Hall  
University of Minnesota  
75 E. River Road  
Minneapolis, MN 55455

Dr. Ronald A. Weitzman  
NPS, Code 54Wz  
Monterey, CA 92152

Major John Welsh  
AFHRL/MOAN  
Brooks AFB, TX 78223

Dr. Rand R. Wilcox  
University of Southern  
California  
Department of Psychology  
Los Angeles, CA 90007

German Military Representative  
ATTN: Wolfgang Wildegrube  
Streitkraefteamt  
D-5300 Bonn 2  
4000 Brandywine Street, NW  
Washington, DC 20016

Dr. Bruce Williams  
Department of Educational  
Psychology  
University of Illinois  
Urbana, IL 61801

Dr. Hilda Wing  
Army Research Institute  
5001 Eisenhower Ave.  
Alexandria, VA 22333

Dr. Martin F. Wiskoff  
Navy Personnel R & D Center  
San Diego, CA 92152

Mr. John H. Wolfe  
Navy Personnel R&D Center  
San Diego, CA 92152

Dr. George Wong  
Biostatistics Laboratory  
Memorial Sloan-Kettering  
Cancer Center  
1275 York Avenue  
New York, NY 10021

Dr. Wendy Yen  
CTB/McGraw Hill  
Del Monte Research Park  
Monterey, CA 93940

DATE  
FILMED  
- 8